INDEX TO VOLUME 49

New names and the final members of new combinations are in **bold face type.**

Abutilon incanum, 246; parvulum, 37
Acacia constricta, 244; Greggii, 244
Acerates lanuginosa, 98; viridiflora, 98
Acer Negundo, 293; rubrum, 165; saccharinum, 293
Achyronychia, 351; Rixfordii, 351
Acrolasia albicaulis, 184; dispersa, 184; humilis, 183; latifolia, 184
Acuan Jamesii, 245
Aecidium **Yuccae**, 195
Aerodiclidium **benense**, 262
Agaricus campanulatus, 232; Carbon, 232; equinus, 232; multifidus, 231; papilionaceus, 232; Sajor-Caju, 230; striatus, 231; varius, 232
Agave, 32; Parryi, 239, 242; utahense, 32
Agropyron, 104; dasystachum, 105; molle, 105; repens, 105; Smithii, 105; tenerum, 105
Albizzia lophanta, 51
Aleurina, 19; elastica, 21; fuscocarpa, 16; marchica, 16; olivacea, 20
Algae in the Chinese classics, References to the, 297
Algae, New British and American species of Lobomonas: a study in morphogenesis of motile, 123
Allionia, 351; Brandegei, 351; nyctaginea, 351; ovata, 351; pumila, 351
Alsophila, 3
Alternaria, 332; Solani, 314, 323, 333–335
Amaranthus paniculatus, 243
Amauria dissecta, 359
Amauriopsis dissecta, 359
Ambrosia, 53
American fossil hepatic, A new, 207
American species of Lobomonas: a study in morphogenesis of motile algae, New British and, 123
Amoeba, 133–135
Amsinckia intermedia, 100; lycopsoides, 100; Menziesii, 100, 239, 247
ANDERSON, FLORA, The development of the flower and embryogeny of Martynia louisiana, 141
Andropogon saccharoides, 241
Anemone, 240; tuberosa, 35; sphenophylla, 239, 243
Anislotus brachycarpus, 240, 245; puberulus, 239, 245; trispermus, 240, 245
Antennaria alpina, 181
Anthericum, 54
Antholyza, 55
Antirrhinum antirrhiniflorum, 248
Aphanolejeunea, 4, 5, 7; sicaefolia, 5
Aplopappus **Gooddingi**, 44; gracilis, 44
Apocynum, 51
Arabis perennans, 244
Aragallus monticola, 179
Arenaria Burkei, 179; congesta, 179; lithophila, 179; subcongesta, 179
Aristida adscensionis, 242; bromoides, 242; divergens, 242; Wrightii, 242
Aristolochia, 51; fimbriata, 51; Sipho, 51, 55
Arizona, Midwinter botanizing in southern, 237
Artemisia albula, 250; Brittonii, 250; pabularis, 102
ARTHUR, JOSEPH CHARLES, New species of Uredineae—XIV, 189
Asclepias eriocarpa, 355; eriocarpa **microcarpa**, 355; linaria, 247; pumila, 98; speciosa, 98; syriaca, 97, 98; verticillata, 98
Ascobolus, magnificus, 304
ASHE, W. W., Notes on trees and shrubs of the southeastern United States, 265
Aspergillus, 328, 376; niger, 375.
Asphodelus, 54
Aspidium simulatum, 63
Asplenium, 6, 8
Astragalus Nuttallianus, 245; parviflorus, 101
Atriplex canescens, 238, 243; hastata, 97; patula, 97; rosea, 243
Audibertia capitata, 37
Auricularia reticulata, 228
Avena, 376
Ayenia pusilla, 246

Baccharis glutinosa, 250; sarothroides, 250
Bahia dealbata, 250; dissecta, 359
Baileya multiradiata, 238, 250
Barbula chloronotus, 239; Manniae, 239

BARTRAM, EDWIN B., Midwinter botanizing in southern Arizona, 237
Basidiomycetes—V, Philippine, 223
Bebbia juncea, 250
Betula, 172; lutea, 172; nigra, 172, 293
Biatorella, 173; resinae, 164, 173
Blechnum, 8; attenuatum, 3, 5, 8
Boletus dichrous, 229; thelephoroides, 229
Bolivia, New species of trees of medical interest from, 259
Botanizing in southern Arizona, Midwinter, 237
Botrytis, 301, 305, 328; cinerea, 305, 314, 317, 325, 327, 335
Bouteloua filiformis, 242; Parryi, 242; Rothrockii, 242
Brachiomonas, 75–77, 79, 80, 82, 84, 85, 87, 89, 92, 125, 131, 132, 135–137; gracilis, 75, 76, 81, 89, 91; **simplex**, 82–85, 89, 91; submarina, 75, 76, 78, 79, 82–84, 89, 91, 92; submarina **obtusa**, 77, 79, 81, 89, 91
Brachiomonas, The phylogeny of the genus, 75
Brachyphyllum, 207
Brachyruscus, 212, 213; **Alleni**, 211–213
Brassica nigra, 244
British and American species of Lobomonas: a study in morphogenesis of motile algae, New, 123
BRITTON, N. L., & ROSE, J. N., Two new genera of Cactaceae, 251
Bromus commutatus, 104; eximius, 176; inermis, 94, 95, 100, 101; japonicus, 104; tectorum, 100; vulgaris eximius, 176
Buchnera elongata, 195
Buechnera, 195
Bulgaria, 21; inquinans, 15; nigrita, 16, 18, 20
Bulgariella nigrita, 16; pulla nigro-olivacea, 16
Bullaria, 190; **Zorniae**, 190

Cactaceae, Two new genera of, 251
California—I, Miscellaneous notes on plants of Southern, 31;—II, 349
Calliandra eriophylla, 239, 244
Callisia, 271
Callistemon lanceolatum, 3, 4
Callitriche palustris, 246
Campanula sibirica, 100, 101
Campelia, 271
Canna, 55
Capsella, 154, 156
Caraguata, 12; Sintenesii, 6, 14
Carbon disulphide, Stimulation of plants by, 375
Carex, 55, 103, 369; alata pulchra, 366; albolutescens, 364, 365, 372, 373; albolutescens cumulata, 366, 373; aquatilis, 176; atherodes, 103; athrostachya, 103; Bicknellii, 369; brevior, 368-370; caespitosa, 177; cumulata, 366; Deweyana, 176; Emoryi, 103; festiva, 177; festucacea, 368–370; foenea, 177, 373; Goodenovii, 177; Heleonastes, 175, 177; Hookeriana, 103; hormathodes, 177, 361; hormathodes invisa, 361; hormathodes Richii, 361, 362; laeviconica, 103; Longii, 373; **Merritt-Fernaldii**, 370; monile, 177; obtusata, 103; Parryana, 103; polygama, 178; praticola, 103; **Richii**, 362; saxatilis, 178; scirpiformis, 103; stenophylla, 103; straminea, 364, 369, 372, 373; straminea Crawei, 369; straminea cumulata, 364, 366; tenera, 103; tenera major, 362; tenera Richii, 361, 362; vesicaria, 177
Carex—XII, Notes on, 361
Carlowrightia arizonica, 249
Carpolithes, 211
Cassia Covesii, 245
Castanea alnifolia, 266, 267; alnifolia floridana, 266; alnifolia pubescens, 266, 267; **Ashei**, 267; dentata, 266, 267; **floridana**, 266, 267; neglecta, 266, 267; pumila, 265-267; pumila Ashei, 267; pumila **Margaretta**, 265, 267; pumila praecox, 267; pumila serotina, 267
Castilleja, 191, 193
Catinella, 15; **elastica**, 16, 21; **nigro-olivacea**, 16, 18-20; olivacea, 16
Catinella, The genus, 15
Caulinia flexilis, 35
Cecidomyia resinicola, 163
Celtis pallida, 243
Centaurea picris, 102
Central America, The genus Costus in, 283
Cephaleuros, 12
Cephalothecium roseum, 320
Cercidium Torreyanum, 245
Cereus giganteus, 247
Chaetochloa macrostachya, 242; viridis, 242
Chamaerhodos erecta, 100
Cheilanthes Covillei, 32; Feei, 32; Lindheimeri, 239, 241; myriophylla, 241; Pringlei, 241; Wrightii, 241
Chenopodium, 95; album, 95–97; ferulatum, 96, 97; leptophyllum, 96, 97; paganum, 96, 243; pratericola, 97
Chilopsis linearis, 249

Chimaphila occidentalis, 355; umbellata, 355
Chinese classics, References to the algae in the, 297
Chlamydomonas, 75, 78, 79, 82, 84, 85, 87–89, 123–125, 127–132; brachyura 80, 84, 85; caudata, 80, 82, 85, 86, 88, 89, 92; metastigma, 128; pulvisculus, 123; reticulata, 127; stellata, 123; subcaudata, 85, 86
Chlorogonium, 128
Chroococcus, 6
Chrysanthemum Leucanthemum, 100
Chrysoma laricifolia, 249
Chrysomyxa, 191
Chrysothamnus formosus, 101; graveolens, 101; nauseosus, 101
Cirsium neomexicanum, 238, 250
Citrus Limonium, 7
Cladosporium, 332
Clastobryum americanum, 5
Claytonia lanceolata **Peirsonii**, 352; umbellata 352
Clematis Drummondii, 244
Clusia havetioides, 5, 6, 8, 9, 13
Cobaea scandens, 53, 55
COCKERELL, T. D. A., A new genus of fossil Liliaceae, 211
Coenogonium, 7
Coleosanthus baccharideus, 249; Coulteri, 249; Wrightii, 249
Collema, 7
Colletotrichum, 332; lagenarium, 323, 325, 329, 330; Lindemuthianum, 333; Schizanthi, 330
Cololejeunea, 4, 6, 7; diaphana, 4–7
Colorado, Two new plants from western, 183
Commelina, 270–272
Commelinaceae, Commelinantia, A new genus of the, 269
Commelinantia, 272; **anomala**, 273–275
Commelinantia, A new genus of the Commelinaceae, 269
Commelyna, 269, 270; anomala, 269; rhodantha, 269
Commicarpus scandens, 243
Condalia spathulata, 246
Copelandia, 232; papilionacea, 232
Corispermum, 97; hyssopifolium, 97; nitidum, 97; villosum, 97
Corticium, 166
Costus, 283, 284; Bakeri, 285, 288, 289; **bracteatus**, 284, 285, 292; comosus, 288; **congestus**, 285, 291, 292; Dinklagei, 286; erythrocoryne, 287; Friedrichsenii, 289; geothyrsus, 287; giganteus, 292; hirsutus, 285, 288; laxus, 285, 292; lima, 285, 288, 289; Malortieanus, 285, 289, 290; maximus, 286, 292; mexicanus, 289; nutans, 285, 291, 292; pictus, 289; pulverulentus, 285, 290; sanguineus, 285, 290; **sepacuitensis**, 284, 286, 292; spicatus, 285, 290, 291; splendens, 285, 292; villosissimus, 284, 287, 288
Costus in Central America, The genus, 283
Coviliea glutinosa, 237, 245
Cracca tenella, 245
Crassina pumila, 250
Crataegus mollis, 100
Crepidotus, 232; pogonatus, 232
Crepis capillaris, 103; occidentalis, 103
Cronartium, 191; coleosporioides, 191; 194; filamentosum, 191, 194; Harknessii, 194; Quercuum, 193; ribicola, 314, 319–321, 323, 329–331, 333, **stalactiformе**, 191, 194
Crossotolejunea, 4–7
Croton, 1
Cryptantha calycosa, 100; gracilis, 39, 40; gracilis **Hillmanii**, 39; Hillmanii, 39, 40; maritima, 38, 39; racemosa, 38; ramosissima, 38, 39; recurvata, 39
Cucurbita, 376; Pepo, 54, 382
Cuscuta, 98, 107; applanata, 109; arvensis, 98; calycina, 98; Choisiana, 107; Coryli, 98; corymbosa grandiflora, 109; **cozumeliensis**, 108; **dentatasquamata**, 107; **durangana**, 109; floribunda, 109; Gronovii, 98, Gronovii curta, 98; macrocephala, 108; partita, 108; pentagona, 98; plattensis, 98; tinctoria, 109; umbellata, 109
Cuscuta from Mexico, Three new species of, 107
Cyanotis, 271
Cyathea, 3
Cylindrocladium scoparium, 320, 328, 330, 334
Cyperus cyrtolepis, 242; diandrus, 103; esculentus, 103
Cystopus candidus, 323, 330, 333

Dacryomyces, 166; deliquescens, 167, 168; hyalinus, 166–168
Daedalea, 225; amanitoides, 225; applanata, 225; aulaxina, 226; flavida, 225; inconcinna, 226; indica, 225; intermedia, 227; lurida, 225; Palisoti, 225; pruinosa, 225, 227; repanda, 225; sanguinea, 224; striata, 231; subconfragosa, 225; **versatilis**, 226
Danaea, 3
Danthonia pinetorum, 175; spicata pinetorum, 175; thermalis, 175

Dasylirion Wheeleri, 239, 242
Dasyochloa pulchella, 242
Daucus pusillus, 247
Desmidium, 4
Development of the flower and embryogeny of Martynia louisiana, The, 141
Diabole, 194; **cubensis**, 194
Diapedium Torreyi, 248
Dicaeoma Polygoni-alpini, 189
Dichorisandra, 271
Didymellina Iridis, 329, 330
Dioscorea, 55; quinquefolia, 51, 55
Diplasiolejeunea pellucida, 4–7
Diplocarpon Rosae, 330
Dipterostemon pauciflorus, 242
Discocactus, 251
Ditaxis neomexicana, 246; sericophylla, 246
DODGE, B. O., A Lachnea with a botryose conidial stage, 301
Dodonaea viscosa, 246
DORAN, WM. L., Effect of external and internal factors on the germination of fungous spores, 313
DOSDALL, LOUISE, Occurrence of the pycnial stage of Puccinia Taraxaci, 235
Draba corrugata, 352; corrugata **saxosa**, 352; cuneifolia, 239, 244; saxosa, 352
Dracocephalum Moldavica, 100; parviflorum, 100; thymiflorum, 100
Drepanolejeunea campanulata, 5–7; crucianella, 7
Dryopetalon, 240, 247; runcinatum, 239, 244
Dryopteris spinulosa, 105; Thelypteris, 105
Dunaliella, 130
DURAND, ELIAS J. The genus Catinella, 15
Dysodia porophylloides, 250
Earliella corrugata, 224; cubensis, 224
Echinocactus, 251; Beguinii, 252; bicolor, 251; conoideus, 252; hexaedrophorus, 251; lophothele, 251; Wislizeni, 247
Echinocereus Fendleri, 247
Echinodorus cordifolius, 35
Effect of external and internal factors on the germination of fungous spores, 313
Elaphoglossum, 3; latifolium, 4, 6, 8–10, 13, 14
Elatine californica, 35
Elmerina, 226; cladophora, 226; vespacea, 226
Encelia farinosa, 240, 250
Endothia parasitica, 320, 330
Ephedra antisyphilitica, 241; trifurcata, 241; viridis, 241
Epicampes ringens, 242
Epidendrum, 7
Epiphyllous plants of certain regions in Jamaica, 1
Eragrostis megastachya, 242
Eremolithia, 351; Rixfordii, 351
Erigeron divergens, 250; grandiflorus, 181
Eriogonum apiculatum, 35; fasciculatum, 351; fasciculatum **flavoviride**, 350; fasciculatum foliolosum, 351; fasciculatum polifolium, 350, 351; nodosum, 350; nodosum **Jaegeri**, 350; ovalifolium, 179; pinetorum, 243; polycladon, 243; Wrightii, 243
Eriophorum polystachyon, 176
Eritrichium racemosum, 38
Erodium cicutarium, 245; texanum, 245
Erucastrum Pollichii, 94
Eschscholtzia mexicana, 244
Eupatorium solidaginifolium, 249
Euphorbia capitellata, 246; Esula, 95; pediculifera, 246; polycarpa, 246
Evolvulus linifolius, 247
Evonymus atropurpureus, 101

Fagus pumila, 267
FARR, CLIFFORD H., Quadripartition by furrowing in Sisyrinchium, 51
Favolus, 228; cucullatus, 227; curtipes, 227; megaloporus, 228; multiplex, 228; princeps, 228; spathulatus, 228
Fendlera rupicola, 244
Festuca Hallii, 104; viridula, 104
Fissidens, 7
Floscopa, 271
Fomes, 172; Bakeri, 172; rimosus, 172; robustus, 172
Forrestia, 271
Foresteria phillyreoides, 247
Fossil, A modern plant, 63
Fossil hepatic, A new American, 207
Fossil Liliaceae, A new genus of, 211
Fouquieria splendens, 246
Fragaria, 53
Franseria deltoidea, 249
Frasera Parryi, 37
Fraxinus attenuata, 247
Freesia, 55
Fritillaria, 56
Fucoides erectus, 207
Fucus, 135
Funalia versatilis, 226
Fungous spores, Effect of external and internal factors on the germination of, 313

Galactia Wrightii, 245

Galium, 358; angustifolium **pinetorum**, 357; californicum, 359; grande, 359; **Hallii**, 358; multiflorum, 358; occidentale, 358, 359; pubens, 359; stellatum, 249, 358
Gayoides crispum, 246
Gentiana humilis, 349; simplex, 349
Germination of fungous spores, Effect of external and internal factors on the, 313
Glacier National Park, Unreported plants from, 175
Gleocapsa, 6
Gloeoporus, 228; candidus, 229; conchoides, 228; dichrous, 229; **reticulatus**, 228
Glomerella rufomaculans, 314, 320, 328, 330
Gnaphalium alpinum, 181; Wrightii, 250
Gonium pectorale, 127, 128
Gouania, 195
GRAFF, PAUL W., Philippine Basidiomycetes—V, 223; Unreported plants from Glacier National Park, 163
Guarea **alborosea**, 263; **Bangii**, 262
Guignardia Bidwellii, 330
Gutierrezia glomerella, 249
Gymnocalycium, 251
Gymnogramme triangularis, 32
Gymnosperma corymbosum, 249
Gymnosporangium clavipes, 323, 330, 331, 334, 335; Juniperi-virginianae, 318, 321–323, 330, 331, 334

Hapalopilus subrubidus, 227
Haplophyton cimicidum, 247
Hartmannia speciosa, 354
HAZEN, TRACY E., New British and American species of Lobomonas: a study in morphogenesis of motile algae, 123; The phylogeny of the genus Brachiomonas, 75
Hedeoma nana, 248; oblongifolia, 248
Hedychium, 3; coronarium, 3, 4
Hedyosmum, 3; arborescens, 4
Hedysarum, 101; cinerascens, 101
Helenium montanum, 250
Helianthus, 53; annuus, 382; giganteus, 101; grosse-serratus, 101; Maximiliani, 101; nitidus, 101; tuberosus, 101
Hepatic, A new American fossil, 207
Herberta, 208; adunca, 208
Hermannia pauciflora, 239, 246
Herniaria cinerea, 35
Hesperonia retrorsa, 243
Heteropogon contortus, 241
Hexagonia, 227, 228; albida, 227; apiaria, 227; Cesati, 227; ciliata, 226; cladophora, 226; **Clemensiae**, 227; cruenta, 224; cucullata, 227; glabra, 226; Koenigii, 227; macrotrema, 226; Molkenboeri, 226; **subrubida**, 227; Taxodii, 227; vespacea, 226
Hibiscus Coulteri, 246; denudatus, 246
Hoffmanseggia densiflora, 245
Holacantha Emoryi, 36
HOLLICK, ARTHUR; HOWE, MARSHALL A., &, A new American fossil hepatic, 207
HOWE, MARSHALL A., & HOLLICK, ARTHUR, A new American fossil hepatic, 207
Humaria, 19; fuscocarpa, 16; marchica, 16, 20; olivacea, 16
Hydrodictyon, 78, 299
Hymenoclea monogyra, 249
Hymenogramme spathulata, 228
Hymenophyllum, 3

Inga vera, 195
Inonotus Clemensiae, 227
Iowa, Quercus lyrata in, 293
Iris, 55
Isocoma Hartwegi, 249
Ixia, 55

Jacquemontia Pringlei, 247
Jamaica, Epiphyllous plants of certain regions in, 1
Jamesoniella autumnalis, 208
Janusia gracilis, 245
Jepsonia Parryi, 36.
JOHNSTON, IVAN M., MUNZ, PHILIP A. &, Miscellaneous notes on plants of Southern California—I, 31;—II, 349
Juglans nigra, 293
Juncus nodosus, 242
Jungermanniopsis, 208; **Cockerellii**, 208
Juniperus scopulorum, 239, 241

Karschia, 15
Koeberlinia spinosa, 245
Koeleria cristata, 242
Krameria glandulosa, 245
Krynitzkia maritima, 38; ramosissima, 38

Lachnea, 301, 305; abundans, 301, 302, 304, 305; cretea, 301, 303, 304
Lachnea with a botryose conidial stage, A, 301
Lactuca sativa, 53; virosa, 103
Laminaria, 299
Lamium amplexicaule, 100
Lappula cenchrusoides, 99; ciliata, 180; cupulata foliosa, 99; floribunda, 99; Lappula, 99; occidentalis, 99; texana homosperma, 99

Laschia, 229; minima, 229; spathulata, 228
Lecanora, 7
Lejeunea flava, 4–7
Lemalis, 18; pulla, 18; pulla nigro-olivascens, 18; rufo-olivacea, 16, 18
Lemna, 103; trisulca, 349
Lentinus, 229, 232; brachatus, 230; Decaisneanus, 230; fusco-purpureus, 229; infundibuliformis, 230; javanicus, 230; Kurzianus, 230; polychrous, 230; praerigidus, 230; Sajor-Caju, 230; setiger, 230; tanghiniae, 230; velutinus, 230; Zeyheri, 230
Lenzites, 225, 231; acuta, 231; applanata, 225; aspera, 226; nivea, 227; Palisoti, 225; pallida, 225; platyphylla, 226, 231; platypoda, 225; repanda, 225; striata, 231
Lepidium densiflorum, 93, 94; Draba, 94; hirsutum, 244; lasiocarpum, 239, 244; ramosissimum, 93, 94
Leptosyne arizonica, 250
Lesquerella Fendleri, 238, 244; purpurea, 239, 244
Liliaceae, A new genus of fossil, 211
Lilium Martagon, 150
Linum spergulinum, 36
Lippia Wrightii, 248
Lobomonas, 89, 123, 124, 128, 130–133, 135–137; Francei, 123–125, 127, 129, 130; **pentogonia**, 124, 125, 127, 129, 139; **rostrata**, 124, 127, 130, 139; stellata, 123
Lobomonas: a study in morphogenesis of motile algae, New British and American species of, 123
Lolium remotum, 104; rigidum, 104; temulentum, 104
Lomatium macrocarpum, 101
Lopadium, 4
Lophocolea, 5–7
Lupinus, 197, 204; affinis, 198, 199, 204; apricus, 204; arvensis, 200, 201; Aschenbornii, 204; Barkeri, 197; bicolor, 204; bilineatus, 197; carnosulus, 204; chihuahuensis, 204; Clarkei, 197; densiflorus, 200, 201; Ehrenbergii, 200, 201; Hartwegi, 197; hirsutulus, 204; leucophyllus, 205; Liebmanni, 199; micranthus, 204; micranthus microphyllus, 204; nanus, 199, 204, 205; niveus, 197, 204, 205; pachylobus, 204; persistens, 204; Pipersmithii, 204; polycarpus, 204; purpureus, 199; ramosissimus, 200, 201; rostratus, 204; sabulosus, 204; strigulosus, 204; succulentus, 197–200; succulentus **Brandegeei**, 203; succulentus **Layneae**, 203; trifidus, 204; umbellatus, 204; vallicola, 204; vallicola apricus, 204
Lupinus succulentus and L. niveus, Studies in the genus Lupinus—VII, 197
Luzula comosa, 349
Lychnis alba, 95
Lycium, 357; Cooperi, 248; Fremontii, 248; parviflorum, 248; Spencerae, 356, 357; Torreyi, 248
Lycopus communis, 100

MACKENZIE, KENNETH KENT, Notes on Carex—XII, 361
Madia glomerata, 102
Magnolia, 51, 55–59
Malacocarpus, 251
Malus elongata, 268; elongata **pubens**, 268; platycarpa **parvula**, 268
Malva borealis, 95; crispa, 95; verticillata, 95
Malvastrum, 353
Mammillaria conoidea, 252; Grahami, 247
Mangifera indica, 1, 5, 10, 13
Marattia, 3
Marchantia Pealei, 207
Marchantites erectus, 207
Marrubium vulgare, 248
Marsilea vestita, 35
Martynia, 141, 150, 154; fragrans, 141, 142; louisiana, 141, 142; lutea, 141
Martynia louisiana, The development of the flower and embryogeny of, 141
Mecardonia peduncularis, 248
Medicago falcata, 100
Melampsoropsis Rhododendri, 191; **roanensis**, 190
Melanconium, 323
Melanospora, 305
Menodora scabra, 247
Merulius fugax, 168, 169; striatus, 231; subaurantiacus, 169
Mesosphaerum Emoryi, 248
Metiola astrina, 12
Metzgeria furcata, 4–7
Mexico, Three new species of Cuscuta from, 107
Microlejeunea, 4, 5
Microstylis monophyllos, 349
Midwinter botanizing in southern Arizona, 237
Mimosa asperata, 194; pigra, 194
Mimulus Langsdorfii, 248
Miscellaneous notes on plants of Southern California—I, 31; —II, 349
Mittenothamnium reptans, 5, 6
Modern plant fossil, A, 63

Mollisia umbonata. 19
Mollugo verticillata, 35
Monbretia, 55
Monoclea Gottschei, 6
Mucronella, 169
Muhlenbergia dumosa, 242; microsperma, 242; Porteri, 242
MUNZ, PHILIP A., & JOHNSTON, IVAN M., Miscellaneous notes on plants of Southern California—I, 31; —II, 349
Mycoidea parasitica, 1
Mycological notes for, 1920, 163
Myoporum serratum, 154
Myosurus cupulatus, 36

Naias, 54; flexilis, 35
Nama humifusum, 356
Nardia, 208
Nectandra, 259, 261; **coto**, 260
Nectriella, 164; resinae, 164
Neolloydia, 251; **Beguinii**, 252; **conoidea** 252
New American fossil hepatic, A, 207
New British and American species of Lobomonas: a study in morphogensis of motile algae, 123
New genus of fossil Liliaceae, A, 211
New genus of the Commelinaceae, Commelinantia, A, 269
New plants from western Colorado, Two, 183
New records and other notes on North Dakota plants, 93
New species of trees of medical interest from Bolivia, 259
New species of Uredineae—XIV, 189
Nicotiana, 55–57, 59; glauca, 248; trigonophylla, 248
Nissolia Schottii, 245
Nolina microcarpa, 242
North Dakota plants, New records and other notes on, 93
Nostoc, 4–6, 299
Notes for 1920, Mycological, 163
Notes on Carex—XII, 361
Notes on North Dakota plants, New records and other, 93
Notes on plants of Southern California, Miscellaneous,—I, 31;—II, 349
Notes on trees and shrubs of the southeastern United States, 265
Notholaena bonariensis, 241; californica, 32; Hookeri, 241; Lemmoni, 239, 241; Parryi, 32; sinuata, 241; sinuata integerrima, 31, 241; tenera, 32
Nuttallia, 183; **marginata**, 183
Nymphaea, 56
Nyssa sylvatica, 166

Occurrence of the pycnial stage of Puccinia Taraxaci, 235
Ocotea, 261; **pseudo-coto**, 261
Oenothera, 354; cardiophylla, 355; cardiophylla **splendens**, 354; speciosa, 354
OLDENBUSCH, CARRIE, Stimulation of plants by carbon disulphide, 375
Olneya tesota, 245
Ophioglossum Alleni, 211
Ophrys monophyllos, 349
Opuntia arborescens, 247; Bigelovii, 238, 247; chlorotica, 247; fulgida, 246; leptocaulis, 247; Toumeyi, 247
Osmunda, 52; palustris aurea, 52; regalis, 52
OSTERHOUT, GEO. E., Two new plants from western Colorado, 183
OVERHOLTS, L. O., Mycological notes for 1920, 163

Panaeolus, 232; campanulatus, 232; papilionaceus, 232
Pandorina, 128
Panicum perlongum, 103
Pannaria rubiginosa, 7
Panus, 232
Papulospora, 305
Parietaria obtusa, 243
Parkinsonia microphylla, 245
Parmelia, 4–6
Parnassia cirrata, 349
Parosela Greggii, 245; Parryi, 245
Parthenium incanum, 250
Patellaria applanata, 16, 19; hirneola, 16; olivacea, 16, 20; pulla, 19; pulla nigro-olivacea, 16, 18; violacea, 16
Patinella applanata, 16, 19; hirneola, 16, 19; olivacea, 16, 19; violacea, 16, 19
Pectocarya linearis, 247
Pediastrum, 131, 132
Pellaea compacta, 32; mucronata, 239, 241
Penicillium glaucum, 334, 375
Penstemon, 43, 189; albomarginatus, 44; antirrhinoides, 43; antirrhinoides **microphyllus**, 43; barbatus, 40; barbatus Torreyi, 40; calcareus, 42; Clevelandi, 41, 357; Clevelandi **connatus**, 357; Clevelandi **Stephensi**, 41, 357; desertorum, 42; Eatoni, 40; fruticosus, 180; Grinnellii, 42; Menziesii, 180; microphyllus, 43; **Munzii**, 40; Palmeri, 41, 42; Palmeri **Grinnellii**, 42; petiolatus, 42; Plummerae, 43; Stephensi, 41; subulatus, 40; superbus, 248
Perezia Wrightii, 249
Peridermium, 193; Cerebrum, 192, 193

filamentosum, 191, 194; Harknessii, 191–193; stalactiforme, 191, 194
Perityle dissecta, 250
Peronospora parasitica, 323; pygmaea, 335
PESSIN, LOUIS J., Epiphyllous plants of certain regions in Jamaica, 1
Pezicula, 19; viridi-atra, 16
Peziza applanata, 18; fuscocarpa, 16, 19, 20; nigro-olivacea, 16, 18, 19; olivacea, 15–17; viridi-atra, 16, 19, 20
Phacelia Franklinii, 180
Phaeopezia, 19; elastica, 21; fuscocarpa, 16, 20; marchica, 16
Phallogaster saccatus, 172
Phaseolus, 376
Philadelphus microphyllus, 244
Philibertella linearis, 247
Philippine Basidiomycetes—V, 223
Phlox, 99; andicola, 99; **bernardina**, 356; dolichantha, 356; Douglasii, 99; Hoodii, 99; Stansburyi brevifolia, 356; superba, 356
Phoradendron californicum, 243; californicum distans **leucocarpum**, 349; macrophyllum, 243
Phragmidium violaceum, 331
Phycopeltis, 3–7, 10; microcystis, 2
Phyllonoma, 212; ruscifolia, 212
Phylloporina epiphylla, 6
Phyllosticta Antirrhini, 323, 330
Phylogeny of the genus Brachiomonas, The, 75
Physalis ixocarpa, 248
Phytophthora infestans, 323, 327, 330, 333
Picea, 191
Pilacre Petersii, 165, 173
Pinnularia, 4–6
Pinus insignis, 193; ponderosa, 164, 193; radiata, 193; Strobus, 167; virginiana, 164
Piper, 3
Pittosporum tobira, 4
Plagiochila, 6
Plasmodiophora Brassicae, 322, 323
Plasmopara viticola, 323, 330, 333, 334
Platanus occidentalis, 293; Wrightii, 244
Platymonas, 80; subcordiformis, 80
Pleurotus, 229; flabellatus, 229
Pluchea, 194; fastigiata, 194; Quitoc, 194
Plumbago scandens, 247
Poa Bigelovii, 242; Cusickii, 176; nevadensis, 176
Podocarpus, 3; coriaceus, 5, 7
Polemonium confertum, 179
Pollia, 271
Polyblepharides, 80, 128
Polygala macradenia, 240, 246
Polygonum alpinum, 189; bistortoides linearifolium, 179; Hydropiper, 97; punctatum, 243; punctatum leptostachyum, 97
Polypodium, 3, 5, 8; Phyllitidis, 7, 8, 10, 14
Polyporus, 228; annosus, 172; apiarius, 227; asper, 223; caeruloporus, 169; Clemensiae, 227; **compactus**, 170, 171, 173; corrugatus, 224; dichrous, 229; Ellisianus, 172; fusco-badius, 224; incanus, 224; indecorus, 224; lacerus, 226; leucoplacus, 223; megaloporus, 228; minimus, 229; nigropurpurascens, 229; paleaceus, 224; pelleporus, 229; scabrosus, 224; subrubidus, 228; tegularis, 224; thelephoroides, 229
Polystichum plaschnichianum, 8
Polystictus, 226; asper, 223; Persoonii, 224; thelephoroides, 229; versatilis, 226
Populus, 243; deltoides, 293; Wislizeni, 243
Poria, 223; leucoplaca, 223
Porophyllum juniciforme, 250
PORTERFIELD, JR. W. M., References to the algae in the Chinese classics, 297
Potentilla argentea, 100
Preissites Wardii, 207
Prosopsis glandulosa, 245
Prunus fasciculata, 356
Pseudopeziza Medicaginis, 317; Trifolii, 317, 330
Psilostrophe Cooperi, 250
Psilotum, 52; triquetrum, 52
Pteromonas, 124, 128, 129, 135; alata, 123
Ptiloria pauciflora, 249
Puccinia, 189; Antirrhini, 316, 323, 330; biocellata, 194; coronata, 321, 323, 332, 334; dispersa, 323; graminis, 321–323, 329, 334; invaginata, 195; Larici, 323; Malvacearum, 314, 322, 323, 329–334; offuscata, 190; Phleipratensis, 323, 330; **Plucheae**, 194; Polygoni-alpini, 189; rubigo-vera, 323; Sorghi, 323; Taraxaci, 235; Zorniae, 190
Puccinia Taraxaci, Occurrence of the pycnial stage of, 235
Pucciniastrum americanum, 190; arcticum americanum, 190
Pyramimonas, 80
Pyrenula, 7
Pyrola picta, 355; umbellata, 355

Quadripartition by furrowing in Sisyrinchium, 51

Quercus, 170; acuminata, 268; alba, 170; aquatica, elongata 268; arizonica, 243; bicolor, 293; **coloradensis**, 268; Emoryi, 293, 243; hypoleuca, 239, 243; lyrata, 293–295; macrocarpa, 268, 293, 294; **nigra elongata**, 268; nigra tridentifera, 268; oblongifolia, 239, 243; stellata, 268; virginiana, 268
Quercus lyrata in Iowa, 293

Refe:ences to the algae in the Chinese classics, 297
Rhamnus tomentella, 246
Rhizina nigro-olivacea, 16, 19
Rhizopus nigricans, 314, 326, 327, 335, 383, 384
Rhododendron, 1, 6, 191; catawbiense, 190; punctatum, 190
Riccardia, 6
ROSE, J. N., BRITTON, N. L., &, Two new genera of Cactaceae, 251
ROUND, EDA M., A modern plant fossil, 63
ROWLEE, W. W., The genus Costus in Central America, 283
Rubus neglectus, 190; occidentalis, 190
Ruellia tuberosa occidentalis, 249
Rumex Britannica, 105; hymenosepalus, 243; paucifolius, 189
RUSBY, H. H., New species of trees of medical interest from Bolivia, 259
Ruscus, 212, 213; hypoglossum, 212; hypophyllum, 213; latifolius, 213
Rutosma purpurea, 246

Salix orbicularis, 178; reticulata, 175, 178, 179; saximontana, 178
Salvia lanceolata, 100; mohavensis, 37; pinguifolia, 248
Salvinia, 211; reticulata, 211, 212
Sambucus mexicana, 249
Samuela, 33
Sarcosoma rufum, 15
Sauvellea, 271
Saxifraga Parryi, 36
Scaphophorus agaricoides, 231
Schizonia vulgaris, 232
Schizophyllum, 231; alneum, 232; commune, 231
Scirpus occidentalis, 103; validus, 103
Scleroma velutinum, 230
Sclerotinia fructigena, 314, 330, 331, 333, 335; Geranii, 301, 302, 305; Ricini, 301
Scopulophila, 351; nitrophiloides, 351; **Rixfordii**, 351
Scrophularia parviflora, 248
Scytonema, 4–7
Sedum Griffithsii, 244
Selaginella arizonica, 241; rupicola, 241
Senecio aureus, 181; crocatus, 181; Lemmoni, 250; manitobensis, 102
Septoria, 323; Gladioli, 330; Lycopersici, 334
SHIMEK, B., Quercus lyrata in Iowa, 293
Shrubs of the southeastern United States, Notes on trees and, 265
Sicyos angulatus, 101
Sideranthus australis, 249; Gooddingi, 44
Silene dichotoma, 95; Douglasii, 179; Fabaria, 95; noctiflora, 95
Silphium, 147
Simmondsia californica, 246
Siphonoglossa longiflora, 249
Sisymbrium Loeselii, 94
Sisyrinchium, 55–57, 59, 195; Bermudiana, 195; Bushii, 56, 61
Sisyrinchium, Quadripartition by furrowing in, 51
SMITH, CHARLES PIPER., Studies in the genus Lupinus—VII. L. succulentus and L. niveus, 197
Solenia fasciculata, 169, 173; villosa, 169
Solidago humilis, 180; Purshii, 180
Sonchus arvensis, 102; uliginosus, 102
Sophia intermedia, 94; ochroleuca, 244
Southern California, Miscellaneous notes on plants of,—I, 31;—II, 349
Sphaeralcea, 353; ambigua, 246, 353; grossulariaefolia, 246; **rosacea**, 353
Sphaerotheca mors-uvae, 318
Spirodela polyrhiza, 103
Sporobolus asper, 104
Staurastrum, 4
Stelis ophioglossoides, 5
Stemodia durantifolia, 248
Stenolobium incisum, 249
Stereum radiatum, 168
Sterigmatocystis, 384, 385; nigra, 376, 383–386
STEVENS, O. A., New records and other notes on North Dakota plants, 93
Sticta, 3, 4
Stimulation of plants by carbon disulphide, 375
Strepanthus arizonicus, 244
Streptolirion, 271
Strigula, 1
Studies in the genus Lupinus—VII. L. succulentus and L. niveus, 197
Study in morphogensis of motile algae, New British and American species of Lobomonas: A, 123
Suckleya Suckleyana, 97
Syntherisma sanguinale, 103

Taxilejeunea, 5–7

THARP, B. C., Commelinantia, a new genus of the Commelinaceae, 269
Thelephora dolosa, 229
Thelocactus, 251; **bicolor**, 251; **hexaedrophorus**, 251; **lophothele**, 251
Thlaspi arvense, 352
Three new species of Cuscuta from Mexico, 107
Thysanocarpus, 240, 247; amplectens, 239, 244
Tilletia foetens, 330
Tinantia, 269, 270, 272; anomala, 270, 273; fugax, 270; Pringlei, 275
Tmesipteris, 211; Alleni, 212
Tortula inermis, 239; ruralis, 239
Tradescantella, 271
Tradescantia, 54, 269–273; anomala, 269, 271–273; Pringlei, 271, 272
Tragia ramosa, 37
Trametes, 223; aspera, 223; corrugata, 224; incana, 223; nitida, 224; paleacea, 224; versatilis, 226
Trapella sinensis, 154
Trees and shrubs of the southeastern United States, Notes on, 265
Trees of medical interest from Bolivia, New species of, 259
Trentepohlia, 3; arborum, 2; aurea, 2, 4, 5, 7
Trichloris fasciculata, 242
Trichomanes, 3, 5
Tricuspis mutica, 32
Triodia mutica, 32
Triticum, 387; sativum, 379
Trixis californica, 249
Tsuga canadensis, 168
Tulasnella, 166; Violae, 166
Two new genera of Cactaceae, 251
Two new plants from western Colorado, 183

Ulmus americana, 293
United States, Notes on trees and shrubs of the southeastern, 265
Unreported plants from Glacier National Park, 163
Uredineae—XIV, New species of, 189
Uredo biocellata, 194; coleosporioides, 194; **cumula**, 195; **curvata**, 195; **nominata**, 194; Plucheae, 194; Zorniae, 190
Uromyces, 189; caryophyllinus, 316, 323; fuscatus, 189; Trifolii, 323
Uromycladium, 194; cubense, 194
Usnea, 4
Ustilago Hordei, 330; Tritici, 330

Vaccaria Vaccaria, 95
Valeriana occidentalis, 180; sylvatica, 180; wyomingensis, 180
Vauquelinia californica, 244
Venturia inaequalis, 314, 317, 318, 320, 325–328, 330–332, 334, 335
Verbena, 148, 150; Wrightii, 248; xutha, 248
Veronica Cusickii, 180; maritima, 98
Vicia, 381, 387; Faba, 380' 381
Villanova dissecta, 359
Vinca rosea, 51
Viola Sheltoni, 353, 354

Wedeliella incarnata, 243
Woodsia scopulina, 31
Woodwardia angustifolia, 63

Xanthium, 101; acerosum, 101; canadense, 101; italicum, 101; pennsylvanicum, 101

Yucca baccata, 33, 34, 239, 242; baccata macrocarpa, 34; elata, 238, 242; glauca, 196; macrocarpa, 34; mohavensis, 33, 34; Schottii, 34
YUNCKER, T. G., Three new species of Cuscuta from Mexico, 107

Zannichellia palustris, 35
Zea Mays, 376
Zebrina, 271
Zythia, 164, 173; resinae, 163, 164, 173